

R. A C T R, BT

Role of olfaction in host plant selection and local adaptation of a polyphagous herbivore, ***Eclyptus*** Sharp

P. R. C. Dodd^{1,*}, M. A. Minor^{1a}, . Wang¹, DJ . Rogers², E. M. Root¹ & . Atreick^{1a}


O: a

A a Y: b (a a a 22.2 c a 4.5 c
 a c a a 24.2 c a 4.5 c a -
) acP a , : P a PaPac-
 P b a a . E P a Y: b
 c cP P a a : e P a
 c cP P a c a c P a acP-
 aP e a c a A a a : P
 a Y: b a a a a
 a P a c a a a Pa aP 400 - / .

I a b P (a Pa P) a P
 P c a a Y: b , :
 a c e a b (4 c a P) a P
 P P aP P ac P aP P
 e a b P P a a . E P e a b
 (4 c a P) c Pa

b a a a 87.3%; $W_{a-} \chi^2_{(1, N = 150)} = 61.86$,
 $P < 0.001$). $M_{a-} (88.3\%) a_{-} a_{-} (86.4\%)$
 $_{,} a_{-}$ ($W_{a-} \chi^2_{(1, N = 150)} = 0.14$,
 $P > 0.05$).

$W_{a-} a_{-} P_{a-} P_{a-} a_{-} P_{a-} a_{-} P_{a-} a_{-} P_{a-}$
 $b_{-} P_{a-} (W_{a-} \chi^2_{(1, N = 36)} = 8.15, P = 0.0043) a_{-}$
 $b_{ac} b_{ac} (W_{a-} \chi^2_{(1, N = 36)} =$



Conclusions

W \dots ab \dots ; \dots \dots c \dots \dots a \dots ; \dots
a \dots a \dots : \dots ca \dots b a \dots E \dots a
 \dots I \dots a \dots a \dots \dots b \dots b \dots
a \dots ; \dots a \dots a \dots a \dots ; b \dots
 \dots \dots a \dots a \dots b \dots ca \dots ; \dots
: \dots a \dots c \dots a \dots . W a \dots
: \dots a \dots a \dots b a a \dots (2
a 3) \dots a \dots a \dots ca
a a \dots . H \dots a \dots a \dots \dots
a \dots (a \dots a \dots Ha \dots c N \dots b \dots
a \dots) a \dots ca a a \dots
a c a \dots a \dots a \dots .
O a \dots a \dots a \dots a \dots a \dots b \dots
a acc \dots c \dots a \dots a \dots a \dots
a \dots a \dots (\dots a \dots) .

Acknowledgements

W \dots a \dots a c a \dots
Ha Ba \dots N Z a a \dots c \dots a
 \dots . PRCD a \dots b L \dots

