

Scientific Research Collaboration (Bibliography): Molecular Zoology

1*, 2, 3, 4, t A. k⁵

1 Department of Morphology, Systematics and Evolutionary Biology, J.F. Blumenbach Institute of Zoology & Anthropology, Georg-August-University, Göttingen, Germany, 2 Department of Anatomy, University of Otago, Dunedin, New Zealand, 3 Department of Ecology, Lincoln University, Canterbury, New Zealand, 4 Independent Researcher, Levin, New Zealand, 5 Ecology Group, Institute of Agriculture and Environment, Massey University, Palmerston North, New Zealand

A t t



1. List of Broscini species used in this study with information regarding authority, distribution (S.I. = South Island New Zealand, N.I. = North Island New Zealand; Ch.Is. = Chatham Islands) and number of individuals employed.

	t t	#	t	A t t
<i>M. alternans</i>	S.I. & Ch.Is.	12		Laporte de Castelnau, 1867
<i>M. alternans hudsoni</i>	The Snares	1		Broun, 1909
<i>M. crenicolle</i>	N.I. & S.I.	9		Laporte de Castelnau, 1867
<i>M. crenaticolle</i>	N.I.	5		Redtenbacher, 1868
<i>M. curvidens</i>	N.I.	1		(Broun, 1915)
<i>M. fulgidum</i> (cf <i>fulgidum</i>)	S.I.	5		Broun, 1881
<i>M. howittii</i>	S.I.	2		Laporte de Castelnau, 1867
<i>M. longicolle</i>	N.I.	1		Broun, 1923
<i>M. lucidum</i>	S.I.	2		Laporte de Castelnau, 1867
<i>M. occiputale</i>	N.I.	3		Broun, 1923
<i>M. cf oconnori</i>	N.I.	4		Broun, 1912
<i>M. oregoides</i>	S.I.	2		(Broun, 1894)
<i>M. rugiceps</i>	S.I.	1		Sharp, 1886
<i>M. sculpturatum</i>	S.I.	4		Blanchard, 1843
<i>M. huttense</i> (cf <i>huttense</i>)	S.I.	3		Broun, 1915
<i>M. simplex</i>	N.I.	4		Laporte de Castelnau, 1867
<i>M. spinifer</i>	N.I.	4		Broun, 1880
<i>M. strictum</i>	S.I.	1		Britton, 1949
<i>M. sulcatum</i>	N.I. & S.I.	1		(Sharp, 1886)
<i>M. validum</i>	N.I.	1		Broun, 1923
<i>M. rectolineatum</i>	S.I.	1		Laporte de Castelnau, 1867
<i>M. politanum</i>	S.I.	1		Broun, 1917
<i>M. impressum</i>	S.I.	1		Laporte de Castelnau, 1867
<i>M. constrictum</i>	S.I.	3		Broun, 1881
<i>M. costellum lewisi</i>	S.I.	1		Broun, 1908
<i>M. costellum obesum</i>	S.I.	1		Townsend, 1965
<i>M. allani</i>	S.I.	1		Fairburn, 1945
<i>M. laterale</i>	S.I.	1		Broun, 1917
<i>M. minax</i>	S.I.	2		Britton, 1949
<i>M. elongatum</i>	S.I.	1		Laporte de Castelnau, 1867
<i>M. metallicum</i>	S.I.	1		Sharp, 1886
<i>M. ducale</i>	S.I.	1		Sharp, 1886
<i>M. morio</i>	S.I.	1		(Laporte de Castelnau, 1867)
<i>M. infimate</i>	S.I.	1		Lewis, 1902
<i>M. punctatum</i>	S.I.	1		(Laporte de Castelnau, 1867)
<i>Meta. moniliferum</i>	S.I.	2		Bates, 1867
<i>Meta. aberrans</i>	S.I.	5		Putzeys, 1868
<i>Meta. tibiale</i>	S.I.	1		(Laporte de Castelnau, 1867)
<i>Brullea antarctica</i>	N.I. & S.I.	1		Laporte de Castelnau, 1867
<i>Bountyia insularis</i>	Bounty Is.	1		Townsend, 1971
<i>O. aereus</i>	S.I.	6		(White, 1846)
<i>O. inaequalis</i>	S.I.	2		(Laporte de Castelnau, 1867)
<i>O. crypticus</i>	S.I.	1		Pawson, 2003
<i>O. septentrionalis</i>	S.I.	2		Pawson, 2003
<i>D. clivoides</i>	S.I.	4		(Laporte de Castelnau, 1867)
<i>D. obtusum</i>	S.I.	2		(Broun, 1886)
<i>D. seclusum</i>	S.I.	1		(Johns, 2007)

doi:10.1371/journal.pone.0086185.t001

2. Cont.

t			
MB 199	<i>Bountya insularis</i>	d	Bounty Is., Proclamation Is.
MB 192	<i>Chylinus ater</i>	d	Australia
MB 13	<i>O. aereus</i>	b	S.I., Otago, Danseys Pass
MB 41	<i>O. aereus</i>	a	S.I., Dunedin, Morrison St.
MB 28	<i>O. aereus</i>	d	S.I., Dunedin, 46 Morrison St.
MB 29	<i>O. aereus</i>	d	S.I., Dunedin, Sandfly Bay
MB 47	<i>O. aereus</i>	d	S.I., Dunedin, Silver Peaks
MB53	<i>O. aereus</i>	d	S.I., L. Onslow, Lammarlows
MB 5	<i>O. inaequalis</i>	d	S.I., Dunedin, Miller Rd.
MB 48	<i>D. clivinooides</i>	b	S.I., Seaward Kaikoura Ra., Tinline Va.
MB 31	<i>D. clivinooides</i>	a	S.I., NW Nelson, Heaphy Track
MB 161	<i>D. obtusum</i>	d	S.I., Fiordland, Kepler Track
MB 162	<i>D. obtusum</i>	c	S.I., Otago, Catlins Coast, Tautuku
MB 159	<i>D. clivinooides</i>	c	S.I., Otago, Kinloch
MB 8	<i>D. clivinooides</i>	d	S.I., Nelson, Cobb Valley
MB 175	<i>D. seclusum</i>	d	S.I., Fiordland, Spey River Valley

(M.= *Mecodema* Meta= *Metaglymma* O.= *Oregus* D.= *Diglymma* S.I.= South Island New Zealand; N.I.= North Island New Zealand; BOP= Bay of Plenty; genes analysed: a=COI, COII, 16S, 18S; b=COI, COII, 16S; c=COI, 18S; d=COI; LUNZ=Lincoln University Entomological Research Museum; P=S.M. Pawson collection); COI sequences from samples of the *Mecodema* group marked with * have previously been deposited in Genbank#(

naturally an increasing interest in how diversification is distributed are predatory *Mecodema* a diverse genus with species distributed throughout the New Zealand mainland from alpine to coastal habitats. In contrast, there is a single species *Mecodema alternans* timing of radiation of *Mecodema* (Blanchard, 1843) carabid beetles the Chatham Islands. The same species occurs in southeast New Zealand near Dunedin. Although *M. alternans* may be better constitutes a prominent species radiation in New Zealand and treated as a species complex [47], no morphological characters presents a good opportunity to explore species level diversification have yet been described that distinguish Chatham Island We utilise the fact that the genus is represented on the Chatham populations from those in mainland New Zealand ([49] & I. Islands, which are located approximately 850 km east of the South Townsend pers. obs.).

In total our sampling comprised 113 specimens, with 88 the formation of this archipelago within the last 4 Myr is *Mecodema* representing 35 described species, and 4 undescribed compelling [3,32,33,34] and corroborated by genetic data for species of the 66 recognized *Mecodema* species (after [45,46] and many taxa (e.g. insects – [18,35], plants – [36,37,38], parakeets <http://www.landcareresearch.co.nz/research/biosystematics/> [39], pigeons – [40,41], cicadas – [42], invertebrates and plants invertebrates/carabid/carabidlist), see Table 1 for details and [43]). In this study the earliest possible establishment of an island authorities. Putative outgroup New Zealand Broscini in our biota (4 Myr) on the Chathams [32] is used as a maximum sample included *Oregus* (Putzeys, 1868) *Diglymma* (Sharp, 1886), possible calibration for estimating the timing of diversification in *Brullea antarctica* (Caporte de Castelneau, 1867) *Metaglymma* (Bates, *Mecodema* Furthermore a substitution rate for Coleoptera [44] is employed to further explore timing of lineage diversification of this beetle genus.


t

t

Sampling


The genus *Mecodema* (Blanchard, 1853) belongs to the tribe Broscini (Carabidae). Broscini has a worldwide distribution but has its main diversity in the southern hemisphere (subfamily Nothobroscinae) [45] and consists of at least 27 genera, comprising about 80 species (see <http://www.landcareresearch.co.nz/research/biosystematics/invertebrates/carabid/carabidlist>) [46,47]. Six endemic genera of Broscini are recognized in New Zealand, but *Mecodema* especially species-rich. Adult *Mecodema* beetles are relatively slow-moving, nocturnal, flightless (with fused elytra), generally active throughout the year, and usually scarce [48]. As with other Carabidae, adults and larvae of the New Zealand taxa

1867), and *Bountya insularis* (Townsend, 1971); plus one representative of Broscini from Australia (*Chylinus atepu* Putzeys, 1868) (Table 2). As many of the species in *Mecodemia* are scarce and therefore difficult to collect we also made use of material from museum collections, and supplemented outgroup sampling with available GenBank sequences (*Oregus septentrionalis* AF466847 & AF466848; *Oregus crypticus* AF54423; *Oregus inaequalis* AF466850 [50]; *Calathus aztecus* GU254333 [51]; *Brososoma relicta* AF012502; *Promecodesps* AF012499; *Creobius eydouxii* AF012498 [52]). The resulting sample represents the geographic and ecological range of *Mecodemia* in New Zealand (Table 2). *Mecodemia* species show examples of allopatric, parapatric and sympatric distribution. Species but not always species groups are limited to particular areas and only few species (*Mg. crenicollis*



2. k f M c a. A) Spatial distribution of samples used in this analysis. Symbols correspond to those in Fig. 2B and code for different clades. B) Analysis of concatenated dataset including mitochondrial COI, COII and 16S plus nuclear 18S. Values at nodes indicate ML bootstrap support returned by analysis using RaxML. Specimen numbers at tips are given as in Table 2. doi:10.1371/journal.pone.0086185.g002

Polymerase chain reactions were performed in 100 volumes and the amplified products then checked on a 1% agarose gel and purified using SAP/EXO1 digest (USB Corporation). Purified PCR products were sequenced using standard protocols for the ABI Prism BigDye Terminator Ready Reaction Kit (Applied Biosystems, Mulgrave, Australia) and run on an ABI Prism 3770 automated sequencer (Applied Biosystems). Sequence identity was so very poor. Although *Dregus* and *Diglymma* represent two of the confirmed by comparison with published data, checked for New Zealand *Nothobroschina* genera considered closest to *Mecodema* [45], it was crucial to verify this relationship within *Broscini* as two other potential outgroup taxa, *Metaglymma* and *Bullea* exist. MrBayes 3.1.2 [59] was used to implement Bayesian analysis with the datasets applying a GTR model with gamma-distributed rate variation across sites and a proportion of invariable sites. Analyses with MrBayes used four independent



3. t f M c a f t . A) COI Bayesian tree generated in BEAST. Numbers on nodes show age estimates based on stratigraphic calibration of 4 Myr. Outgroup taxa were the same as used in the outgroup test (Fig. 1). Grey branches indicate lineages present in North Island New Zealand, and black branches indicate South Island lineages. Tip symbols correspond to clade and location identifiers in Fig. 2. Coloured symbols match symbols in Fig. 3B, highlighting disjunct lineages in southern North Island and northern South Island. Asterisks indicate age estimates between N.I. and S.I. lineages (* = 1.60 Myr, ** = 1.69 Myr, *** = 2.03 Myr). B) Reconstruction of the paleogeographic environment in lower North Island, New Zealand ca. 3 million years ago, green areas indicating likely land above sea level during this time (modified from [28]). Black outlines indicate present day New Zealand land area with coloured symbols corresponding to those in Fig. 3A, showing the present sampling

4. BEAST time estimates based on stratigraphic and COI substitution rate of Coleoptera [44] calibration.

	t	t	(5%)	t	t	t	t	t	t	(5%)	t	t	t
A	4							0.2	(0.0	0.46)			
B	1.24	(0.5–2.09)						0.07	(0.02–0.15)				
C	8.38	(5.29–12.06)						0.52	(0.22–0.9)				
D	6.39	(3.97–9.41)						0.4	(0.18–0.72)				

Markov Chain Monte Carlo (MCMC) runs for ten million generations with a burn-in of 10% and a tree sampling frequency of 1000. Results were checked for convergence. Resulting posterior probabilities on the nodes were recorded.

To examine the species phylogeny of *Mecodema* group in New Zealand we employed all four genes (three mitochondrial and one nuclear) with a subset of 50 taxa (44 ingroup and 6 outgroup samples). The outgroup sampling was chosen after consideration of the results from the prior outgroup analyses. All taxa with data missing for no more than one of four genes were included in the phylogenetic analysis (Table 2). Partition-homogeneity tests (PHITS [60]) were implemented in PAUP*4.0b10 [61] with 500 replicates for the combination of the gene regions to detect significant heterogeneity among the data sets.

MrBayes 3.1.2 [59] was then used to implement Bayesian analysis with the concatenated dataset, applying a GTR model with gamma-distributed rate variation across sites and a proportion of invariable sites. The same model was applied to the partitions with rates and nucleotide frequencies for each gene unlinked. Analyses with MrBayes used four independent Markov Chain Monte Carlo (MCMC) runs for two million generations with a burn-in of 25% and a tree sampling frequency of 1000. Resulting posterior probabilities on the nodes were recorded. The same data were subjected to Maximum Likelihood analysis with bootstrap resampling incorporating a GTR model with gamma-distributed rate variation. ML analysis used RaxML [62] implemented via the CIPRES portal [63]. The data were partitioned by gene (COI, COII, 16S, 18S) and bootstrap resampling was halted by RaxML

Molecular dating

As fossil remains of *Mecodema* that could provide information for calibrating a molecular clock have not been found, yet, we had to rely on geological information and a substitution rate calculated for Coleoptera COI [44] for calibration. In order to gauge the timing and extent of species radiation within New Zealand COI sequences were obtained for additional specimens in addition to previous analyses (Table 2). In some cases, this drew upon museum specimens to further assess the stability of our inferences about the distribution of diversity and timing of radiation in this beetle group.

In total 113 unique COI sequences were used for molecular dating in *Mecodema* (Table 2). To obtain estimates for the maximum age of lineage formation within the genus *Mecodema* we used this dataset of COI with two different calibration strategies. First we employed a stratigraphic calibration to estimate divergence times using the split between Chatham Island

and its closest relative on mainland New Zealand. We assumed a normal distribution for the age around a calibration value of 4 Myr, derived from the maximum age for the Chatham Islands land surface [3,33], assuming that colonization was most likely sooner after emergence of the islands than later. Alternatively, to capture the minimum likely diversification dates we also calibrated the COI dataset with the substitution rate estimated by Pons et al. [44] for Coleoptera COI. This included a normally distributed prior on the substitution rate of 0.08606 subst/site/myrs/l, and a 95% HPD interval from 0.0253–0.147 subst/site/myrs/l as an approximation to the posterior distribution provided by Pons et al. [44]. This rate obtained from analysis of numerous beetle taxa is amongst the highest estimated for any animal gene, and other rates obtained for particular beetle lineages are much slower (e.g. 0.0211 subst/site/myrs/l [64], 0.02 subst/site/myrs/l [65]). Even these rates are nearly twice the widely employed 0.0115 subst/site/myrs/l estimate of Brower [66]. Age estimation for both datasets and both calibration strategies were conducted under the assumption of a strict molecular clock as well as assuming a relaxed molecular clock with a lognormal distribution of rates along the phylogenies [67]. The fit of both priors was compared using Bayes Factors. For all datasets and calibration strategies the relaxed lognormal clock fitted the data decisively better than the strict clock [68].

We used the software BEAUTI 1.4.8 [69] and BEAST 1.7.5 [70] for molecular dating with the given calibrations. All analyses were conducted with a Birth-Death tree prior and a random starting tree under the GTR+I+C model of nucleotide substitution. The MCMC was run for 50–100 million generations, sampling every 5000–10,000th step after a discarded burn-in of 5–10 million steps. Each analysis was run at least two times. The program Tracer 1.4 [71] was used to summarize posterior distributions of all parameters in question, to verify convergence of the MCMC and to estimate Effective Sample Sizes (ESS). If the effective sample size was less than 200, a third MCMC run was conducted for the respective analysis. After convergence of the MCMC was confirmed, the posterior distributions of all parameters were estimated from the combined posterior distributions of all runs conducted for each analysis. The program FigTree 1.4.0 [69] was used to visualize the reconstructed phylogenies.

Three widely used mitochondrial gene regions were employed among species of *Mecodema* with a maximum ML-distance of 0.0179 in COII (COI: 0.0161, 16S: 0.053). Lower values for 16S

compared to COI and COII reflect the comparatively low proportion of variable sites in this gene (16.9%).

Separate analyses of COI and 18S DNA sequences from 41 specimens of *Mecodema* and 6 outgroup taxa (Table 2) resulted in similar topologies even though sequence variation in 18S was low. These analyses confirm *Oregus* and *Diglymma* as the sister group to *Mecodema* and revealed the placement of *Metaglymma* and *Brullea* within the *Mecodema* radiation (Figs. 1A & B).

The alignment of data from four gene regions comprising 50 specimens sampled across New Zealand (including 6 outgroup specimens – Table 2, Fig. 2) was 3114 bp long in total. All three mitochondrial genes displayed the average insect A-T content of about 75%. The partition homogeneity test (PHT) revealed no significant heterogeneity of lineage partitioning among the data sets ($p=0.866$), suggesting their concatenation was appropriate. The GTR+I+C model of nucleotide substitution was identified as the best fitting model by the hLRT and the AIC as implemented in Modeltest 3.5 [72].

Bayesian and ML analysis of the concatenated dataset supported a single topology with the same groupings and branching order and well-supported nodes (Fig. 2). *Mecodema* was confirmed as paraphyletic with respect to *Metaglymma* and *Brullea* as these fall inside the *Mecodema* complex throughout all datasets and analyses in this study. We note that in all cases *Metaglymma* and *Brullea* are placed within the *M. curvidens/origo* clade. This phylogenetic position contradicts the current taxonomic classification and needs to be addressed further in the future. Clades revealed in this analysis comprise species that are, in many cases,

by the setting of priors. In this study the inferred mitochondrial divergence times based on the fast COI rate obtained for Coleoptera [44] are consistently substantially younger than would be expected for a highly diversified genus even in the relatively youthful landscape of New Zealand.

Despite a perception that New Zealand is an ancient land mass

demonstrating the population genetic and ecological mechanisms of diversification (e.g. [89]). Intergenerational population genetics of large invertebrate populations will provide the basis of exciting research.

Increasingly, the fields of species phylogenetics and population phylogeography are merging as it becomes easier to generate appropriate DNA data, and the focus in taxonomy is shifted towards an evolutionary paradigm (e.g. [90]). Teasing apart the interaction of abiotic and genetic processes on population subdivision remains challenging but one taxon group that will provide helpful insight, and it is already evident that *Mecodema* is an impressive example of recent species radiation in the New Zealand fauna. In recent years, synthesis of phylogenetic, ecological and taxonomic evidence has indicated that the biology of New Zealand is primarily the story of recent adaptation and speciation [8,9,19]. Understanding properly how our view of “recent” geological time relates to the

A k t
We would like to thank Steve Pawson and Peter Johns for contributing specimens to this study. Mary Morgan-Richards and Frank Wieland gave valuable comments on earlier drafts of the manuscript. The Department of Conservation assisted with permitting.

A t t t
Conceived and designed the experiments: JG SAT. Performed the experiments: JG. Analyzed the data: JG MK SAT. Contributed reagents/materials/analysis tools: JG MK RME JIT SAT. Wrote the paper: JG SAT.

f

1. Daugherty CH, Gibbs GW, Hitchmough RA (1993) Mega-island or micro-continent? New Zealand and its fauna. *Trends Ecol Evol* 8: 437–442.
2. Trewick SA, Morgan-Richards M (2009) New Zealand Biology. In *Encyclopedia of Islands*. (R. G. . Gillespie and D. A. . Clague, eds.) University of California Press (pp. 665–673).

45. Roig-Juñent S (2000) The subtribes and genera of the tribe Broscini (Coleoptera: Carabidae): Cladistic analysis, taxonomic treatment, and biogeographical consideration. *Bull Am Mus Nat Hist* 255.
46. Britton EB (1949) The Carabidae (Coleoptera) of New Zealand, Part III – A revision of the tribe Broscini. *Trans R Soc N Z* 77: 533–581.
47. Laroche A, Larivière M-C (2001) Carabidae (Insecta: Coleoptera): catalogue. *Fauna of New Zealand*, Number 43 (pp. 44–61). Lincoln, Canterbury, New Zealand, Manaaki Whenua Press.
48. Hutchison MAS (2001) Habitat use, seasonality and ecology of carabid beetles (Coleoptera: Carabidae) in native forest remnants, North Island, New Zealand. Master thesis, Palmerston North, Massey University, New Zealand.
49. Townsend JI (1971) Entomology of the Aucklands and other islands south of New Zealand: Coleoptera: Carabidae: Broscini. *Pacific Island Monographs* 27: 178–180.