

Heteroplasmy of Mitochondrial DNA in the Ophiuroid *Astrobrachion constrictum*

D. J. Steel, S. A. Trewick, and G. P. Wallis

We demonstrate the presence of mitochondrial heteroplasmy for the cytochrome oxidase I (COI) gene of the brittle star (*Astrobrachion constrictum*). One of the 117 individuals analyzed contained two distinct single-strand conformation polymorphism (SSCP) haplotypes differing by two substitutions; another showed sequence evidence for heteroplasmy. We used polymerase chain reaction (PCR) cloning, SSCP, and sequencing of a 480 bp region of the 5' end of COI to isolate and characterize these haplotypes. This is the first properly substantiated case of heteroplasmy in an echinoderm species and may have arisen from paternal leakage.

Mitochondrial DNA (mtDNA) has become a powerful tool for assessing relationships among individuals, populations, and species of animals (Avice 1994). As the number of studies using this genome increases, knowledge of the genetics of the genome itself is also increasing. Two of the more surprising discoveries have been the extent of heteroplasmy in animal populations (Lunt et al. 1998) and cases of biparental inheritance of the genome (Gyllenstein et al. 1991; Hoeh et al. 1991). Heteroplasmy, the occurrence of more than one type of mtDNA in the same organism, can arise either from mutation of the genome within the individual, heteroplasmy of the original oocyte, or from biparental inheritance. Most published examples of heteroplasmy involve a variation in the number of repeats within the control region of the genome (Lunt et al.

1998). Although the control region is non-coding, it probably contains sequences that initiate replication and transcription (Clayton 1982). In echinoids and vertebrates, the displacement loop (d-loop) structure evidences replication (Matsumoto et al. 1974). The length of repeats found in this region ranges from small microsatellite-like repeats (Wenink et al. 1994) to large repeats of 1100 bp (Wallis 1987). Length heteroplasmy is generally explained by slipped-strand mispairing during replication (Densmore et al. 1985), and high frequencies of length heteroplasmic individuals can occur in some species (Lunt et al. 1998). In a few cases observed heteroplasmy has been attributed to biparental inheritance (Kondo et al. 1990; Magoulas and Zouros 1993). Pa-

Figure 1. (A) SSCP gel of COI PCR products from heteroplasmic individual, ER11 (haplotypes A/D), with homoplasmic individuals either side (haplotypes A and D). Other haplotypes found in the Fiordland pop-

A GGAAATTGAT TAGTTCCTT AATGATAGGA GCGCCAGACA TGGCATTCCC TCGAATGAAT AACATGAGAT




Figure 2. COI nucleotide and amino acid sequences for the haplotypes observed in the two heteroplasmic individuals.

replications, or there has been paternal leakage. For the first to occur, heteroplasmy would have to have persisted long enough in the same lineage to evolve two independent mutations.

From studies on *Drosophila* (Solignac et al. 1983) and crickets (Rand and Harrison 1986) it has been suggested that fixation is complete within a few hundred generations. This is a short time for two mutations to have evolved before sorting out into homoplasmic lineages, but without exact knowledge of the sorting out rates and mutation rates within COI of echinoderms we cannot discount this possibility. The second explanation, paternal leakage, seems more plausible. Experiments de-

signed to detect low levels of paternal leakage through repeated backcrossing have shown partial paternal mitochondrial inheritance in *Drosophila* (Kondo et al. 1990) and mice (Gyllensten et al. 1991). These studies suggested that the observed heteroplasmy may be a result of reduced compatibility between egg and sperm due to the use of hybrid strains. However, heteroplasmy attributed to paternal input has been observed in natural




Figure 2. Dental arches of the X trisomy bitch: (A) upper dental arch, (B) lower dental arch. The missing premolar teeth are indicated by arrows.

curved on both sides instead of being straight. The uterine cavity was filled with a small amount of gray mucus. Ovaries were of normal shape and size.

In the bitch's mouth it was found that some premolar and molar teeth were missing in the upper and lower dental arches

three of the eggshell weight estimates. Partial correlation analyses in the two most frequent genotypic classes, *PstI*(1/1) and *PstI*(1/2), revealed the presence of a regulatory loop between feed consumption, body weight, egg weight, and the rate of egg laying. Several aspects of this regulatory loop were different between the two genotypic classes. In particular, for the *PstI*(1/1) genotype, feed consumption was positively associated with egg weight, while there was no significant association for the *PstI*(1/2) genotype. Further, the degree of association of body weight with egg weight decreased with age in the genotypic class *PstI*(1/2), while it was constant for the *PstI*(1/1) genotype. The results indicated that the marker in the IGF-I gene was not only associated with changes in some trait means, but also with changes in the stability of the coordination between*




Figure 1. Flow chart of trait measurements. AFE, age at first egg; HDR, rate of egg laying; EWT, egg weight; SPG, specific gravity; HBWT, housing body weight; MBWT, mature body weight; FBWT, final body weight; FC, feed consumption.

Table 1. Association between IGF-I genotypes and traits means

Traits ^a	Significance		IGF-I genotype			Orthogonal contrasts ^b (P values)	
	Kruskal-Wallis	ANOVA	<i>PstI</i> (2/2)	<i>PstI</i> (1/2)	<i>PstI</i> (1/1)	Linear (additive)	Quadratic (dominant)
			(N 5 12-14) ^c	(N 5 83-97)	(N 5 208-248)		
AFE (d)	0.17	0.43	167	167	166	0.808	0.351
HBWT (g)	0.145	0.083	1242	1267	1297	0.160	0.884
MBWT	0.192	0.192	1715	1729	1776	0.320	0.656
FBWT	0.247	0.216	1689	1706	1745	0.327	0.711
HDR1 (%)	0.269	0.140	85.8	85.7	83.7	0.415	0.624
HDR2	0.256	0.365	74.3	70.5	69.2	0.252	0.750
HDR3	0.439	0.576	55.1	56.7	54.3	0.926	0.460
EWT1 (g)	0.052	0.045	50.3 ^d	52.8 ^e	52.5 ^e	0.006	0.023

Table 3. Analysis of variance of factor means^a

Genotype	<i>N</i>	Factor means ^b				
		F1 (BWT, FC)	F2 (HDR)	F3 (SPG)	F4 (EWT)	F5 ^c (HBWT, AFE)
<i>Psfl</i> (2/2)	10	20.296	20.1132			

genotypes. Additional contributions in decreasing order are the factor which represented the rates of egg laying, and the factor which represented body weight plus feed consumption (Table 3).

Partial correlation analyses (i.e., correlation between two traits corrected for the influence of the remaining traits) indicated the presence of a regulatory loop which coordinates body weight, feed consumption, egg weight, and the rate of laying (Figure 3). Thus in the *Psd(1/1)* genotypic class, heavy chickens tended to lay

1995. Insulin-like growth factor I and parathyroid hormone effects on the growth of fetal rat metatarsal bones cultured in serum-free medium. *Biol Neonate* 68:368–376

Davoren JB, Hsueh JW, and Li CH, 1985. Somatomedin C augments FSH-induced differentiation of cultured rat granulosa cells. *Am J Physiol* 249:E26–E33.

Donoghue DJ, Campbell RM, and Scanes CG, 1990. Effect of biosynthetic chicken growth hormone on egg production in White Leghorn hens. *Poult Sci* 69:1818–1821.

Feng XP, Kuhnlein U, Aggrey SE, Gavora JS, and Zadworny D, 1997. Trait association of genetic markers in the growth hormone and growth hormone receptor genes in a white leghorn strain. *Poult Sci* 76:1770–1775.

Feng XP, Kuhnlein U, Fairfull WR, Aggrey SE, and Zadworny D, 1998. Association of a genetic marker near the growth hormone receptor gene with juvenile body weight in chickens. *J Hered* 89:355–359.

Gavora JS, Kuhnlein U, Crittenden LB, Spencer JL, and Sabour MP, 1991. Endogenous viral genes: association with reduced egg production rate and egg size in White Leghorns. *Poult Sci* 70:618–623.

Goddard C, Wilkie R, and Dunn IC, 1988. The relationship between insulin-like growth factor, growth hormone, thyroid hormone and insulin in chickens selected for growth. *Domest Anim Endocrinol* 5:165–176.

Gowe RS and Fairfull RW, 1990. Genetic controls in selection. In: *Poultry breeding and genetics* (Crawford RD, ed). Amsterdam: Elsevier; 935–954.

Gowe RS, Fairfull RW, McMillan L, and Schmidt GS, 1993. A strategy for maintaining high fertility and hatchability in a multiple-trait egg stock selection program. *Poult Sci* 72:1433–1448.

Harvey S and Hull KL, 1997. Growth hormone: A paracrine growth factor? *Endocrine* 7:267–279.

Isaksson OGP, Eden S, and Jansson JO, 1985. Mode of action of pituitary growth hormone on target cells. *Annu Rev Physiol* 47:483–499.

Johnson RW, Arkins S, Dantzer R, and Kelley KW, 1997. Hormones, lymphopoietic cytokines and the neuroimmune axis. *Comp Biochem Physiol A Physiol* 116:183–201.

Kajimoto Y and Rotwein P, 1991. Structure of the chicken insulin-like growth factor-I gene reveals conserved promoter elements. *J Biol Chem* 266:9724–9731.

Klein S, Morrice DR, Sang H, Crittenden LB, and Burt DW, 1996. Genetic and physical mapping of the chicken IGF1 gene to chromosome a and conservation of synteny with other vertebrate genomes. *J Hered* 87:10–14.

Kocamis H, Kirkpatrick-Keller DC, Klandorf H, and Killefer J, 1998. *In ovo* administration of recombinant human insulin-like growth factor-I alters postnatal growth and development of broiler chicken. *Poult Sci* 77:1913–1919.

Kuhnlein U, Ni L, Weigend S, Gavora JS, Fairfull W, and Zadworny D, 1997. DNA polymorphisms in the chicken growth hormone gene: response to selection for disease resistance and association with egg production. *Anim Genet* 28:116–123.

Li S, Aggrey SE, Zadworny D, Fairfull W, and Kuhnlein U, 1998. Evidence for a genetic variation in the mitochondrial genome affecting traits in White Leghorn chickens. *J Hered* 89:222–226.

McCann-Levorske LM, Radecki SV, Donoghue DJ, Malamed S, Foster DN, and Scanes CG, 1993. Ontogeny of pituitary growth hormone and growth hormone mRNA in the chicken. *Proc Soc Exp Biol Med* 202:109–113.

McMurtry JP, Francis GL, Upton FZ, Rosselot G, and Brocht DM, 1994. Developmental changes in chicken and turkey insulin-like growth factor-I (IGF-I): studies with a homologous radioimmunoassay for chicken IGF-I. *J Endocrinol* 142:225–234.

Monaco MH and Donovan SM, 1997. Insulin-like growth factor-I increases *in vivo* skeletal muscle and mammary

α -amino [^{14}C] isobutyric acid accumulation in food restricted lactating rats. *Nutr Res* 17:1143–1154

Ohlsson C, Bengtsson B, Isaksson OGP, Andreassen TT, and Słotweg MC, 1998. Growth hormone and bone. *Endocr Rev* 19:55–79.

Radecki SV and Scanes CG, 1997. The hypothalamo-pituitary growth hormone-insulin-like growth factor axis: coupling and uncoupling. Perspectives in avian endocrinology (Harvey S and Etches R, eds). Bristol: Journal of Endocrinology Ltd; 119–130.

Scanes CG, Dunnington EA, Buonomo FC, Donoghue DJ, and Siegel PB, 1989. Plasma concentrations of insulin-like growth factors-I and -II in dwarf and normal chickens of high and low weight selected lines. *Growth Dev Aging* 53:151–157.

Schoenle E, Zapf J, Humbel RE, and Froesch ER, 1982. Insulin like growth factor stimulates growth in hypophysectomised rats. *Nature* 296:252.

Soares JH Jr, 1984. (bEndocrinol-o-pi-)wth and no 1[(MInsalr)2rCnnsalrocht 2tedton]TJT* [(phy4cting-338(ER,-1.714 engtsson)

Table 1. Species and DNA sequence used

Species pair	bp	p _b	Number unique	p _w
Genus <i>Haemonchus</i>	459	0.152		
<i>H. contortus</i>			37 (50)	0.025
<i>H. placei</i>			31 (40)	0.019
Genus <i>Teladorsagia</i>	390	0.134		
<i>T. circumcincta</i>			39 (40)	0.023
<i>T. boreoarcticus</i>			8 (11)	0.005
Genus <i>Heteror</i>				

DNA such as this because there are effectively fewer possible character states, so sites saturate quickly (e.g., Brower and DeSalle 1998; Wolfe and Sharp 1993). Unfortunately, how one would accurately correct these tables for multiple hits is not obvious. Any post hoc correction to the tables will need to incorporate a realistic model of the mode of substitution, which in nematode DNA is extremely biased and is not fit by any of the standard models (Blouin et al. 1998).

Further research is needed to determine if selection really acts differently in nematode mtDNA, or if the MK test results are purely an artifact of silent-site saturation. The simplest test would be to find pairs of nematode species that have not diverged as much as these species pairs. Although

solved in detail but is regarded as a polyketide synthesis (Cardani et al. 1973). This pathway implies the presence of multifunctional proteins that biosynthesize (part of) the substance through metabolic channeling (Luckner 1990).

After experimental application of *Paederus* hemolymph to human skin, no reactions were observed by Ito (1934) using *P. poweri* and by de Leon (1952) using *P. fuscipes*, two species which were shown by other authors to cause dermatitis (Frank and Kanamitsu 1987). Such a negative result was attributed to immunization of the test person (Théodoridès 1952). Recent chemical analysis of *P. riparius* and *P. fuscipes* (Kellner and Dettner 1995), however, indicate that contradicting evidence using the same species is a real phenomenon due to pederin polymorphism. In both species studied, most of the females accumulate pederin and transfer it into their eggs, whereas some females are obviously unable to biosynthesize the substance and lay eggs without pederin. The former are concisely called (1) females, the latter (2) females. Like the (2) females, larvae and males do not increase their pederin content by themselves but sequester the substance received maternally or consumed if given access to conspecifics.

Polymorphism for a defensive compound is known in great detail from an example in plants: *Trifolium repens* has a cyanogenic and an acyanogenic morph which differ in mollusk acceptability (Dirzo and Harper 1982). Cyanogenesis, the production of HCN, has long been known to be dependent on the presence of cyanogenic glucosides and a specific β -glucosidase (Jones 1972). It is widely accepted that the cyanogenic polymorphism is controlled by alleles of two loci (Hughes 1991): Alleles at locus *Ac* determine the presence or absence of two cyanogenic glucosides, linamarin and lotaustralin, while alleles at locus *Li* regulate the presence or absence of linamarase, a β -glucosidase that hydrolyzes linamarin and lotaustralin. The loci segregate independently according to Mendelian ratios.

After discovering the pederin polymorphism in *Paederus*, Kellner and Dettner (1995) hypothesized that this polymorphism might also be explained by genetic differences. Heterozygous (1) females could then produce homozygous (2) females, which were surmised because some females descended from (1) females had not accumulated pederin when they were analyzed several months after ima-

ginal eclosion. As pederin is present in the hemolymph all the time and not only after liberation by an enzyme after predation as in cyanogenesis, one locus could suffice for the distinction between (1) and (2) females. Analyzing the progeny of known specimens reared in the laboratory, this study aims at finding evidence for or against such a genetic basis of pederin polymorphism.

Materials and Methods

Beetles

Adult rove beetles (*Paederus riparius*) are found in central Europe mainly in spring and autumn (Horion 1965). The beetles reproduce in spring and imagoes of the new generation hibernate (Boháč 1985). Therefore beetles collected in northeastern Bavaria, Germany, from autumn 1992 to spring 1996 were grouped according to their expected season of reproduction, that is, the 1992 autumnal catch was combined with the beetles collected in spring 1993 under the label 1993 and so forth. Nine sites in northeastern Bavaria were visited, some repeatedly, to collect *P. riparius*: two sites in 1993, four in 1994, six in 1995, and two in 1996. The sites lie up to 100 km apart.

In the laboratory the beetles were isolated according to sex and site. Those collected in autumn had to be hibernated artificially by placing them for at least 3 months in a dark climate chamber at 6°C. After that period or upon collection in spring, pairs were founded and kept separately as described by Kellner and Dettner (1995) in order to obtain eggs of particular females. The eggs were taken out of the breeding cages three times per week and the larvae reared on moist absorbent paper in 24-cell wells (1.7 cm diameter of the wells). Frozen *Drosophila melanogaster* flies were supplied twice a day. One feeding during each larval stage (first and second stadium) consisted of a piece of either a *Tenebrio molitor* larva or a *Calliphora* pupa, which reduces larval mortality to about 22% (Kellner 1998).

The first-generation laboratory-reared imagoes were kept singly in petri dishes (9 cm diameter) with moist absorbent paper where they were fed with live *D. melanogaster* (strain vg). After artificial hiber-




Table 1. (1) and (2) females in the progeny of 17 *P. riparius* (1) females

Figure 1. Percentage of (1) females (with 95% confidence interval) in *P. riparius* samples collected from several sites in northeastern Bavaria during four consecutive years. The females were caught during the preceding autumn or in spring of the respective years and laid eggs in the laboratory containing or lacking pederin.




Figure 3. Percentage of (1) and (2) females in successive laboratory generations of *P. riparius*. Collected (2) females (P) are of unknown descent (?).

Table 3. Amount of pederin determined in *P. riparius* (1) females

(1) Females	<i>n</i>	Pederin (mg)	
		Mean ± SEM	Range
Collected	29	10.8 ± 0.62	4.5-16.8
Kept in laboratory	16	12.5 ± 1.05	5.3-21.0
Laboratory reared	39	14.7 ± 0.92	4.5-24.8
Dead (1)			

deterred by pederin (Kellner and Dettner 1996) cannot be blamed for that because they reject all progeny of (1) females, that means future (2) females as well. Abiotic factors such as a distinct hibernation rate between (1) and (2) females can be ruled out, as the females collected in autumn and hibernated artificially gave no indication of such a factor's importance.

Regarding the data discussed, it is clear that the initial hypothesis is not supported because the ability to biosynthesize pederin cannot be inherited from the father and furthermore no Mendelian proportions are found in the progeny of (1) mothers. The sudden drop of the percentage of (1) females in F_1

vided support for specific differentiation




Figure 1. Composite G-banded (on the left) and C-banded (on the right) karyotypes of a male *P. slevini* (FN 556). Chromosomes 1, 3, 9, 22, 23, and the sex pair are submetacentric or metacentric; the remaining chromosomes are acrocentric. Heterochromatin is restricted to the centromeric region and the sex pair.



Figure 2. Composite G-banded (on the left) and C-banded (on the right) karyotypes of a male *P. sejugis* (FN 5 76). With the exception of chromosome 13, all individuals had indistinguishable karyotypes. Karyotypes from specimens collected on Isla San Diego (the pair on the left) exhibited an alternate form of the submetacentric chromosome 13, designated 13B9. Karyotypes from individuals from Isla Santa Cruz (the pair on the right) have a heterochromatic addition to that chromosome, 13B91. For all individuals, noncentromeric heterochromatin is present on chromosomes 11, 18, and 21.

ated with the *P. maniculatus* species group. Excluding *P. slevini*, the taxa in the *P. maniculatus* species group share indistinguishable biarmed inverted and derived conditions of chromosomes 2, 3, 9, and 20, with the latter character state occurring only in this group and convergently in one cytotype of *P. leucopus* (Stangl 1986). The karyotype of *P. slevini* exhibits acrocentric conditions of both chromosomes 2 and 20. Further, the composite array of chromosomal conformations in the karyotype of *P. slevini* is unique among all reported G-banded karyotypes of deer mice; no other species of *Peromyscus* is known to exhibit the combination of an acrocentric chromosome 2 and biarmed chromosomes 3 and 9.

Although comparisons of the G-banded karyotypes among species of *Peromyscus* do not yield an unambiguous species-group association of *P. slevini*, these data do provide initial hypotheses for studies designed to resolve the phylogenetic position of this species. The karyotype of *P. slevini* is most similar to those that generally characterize taxa in the *P. boylii* and *P. mexicanus* species groups. From the FN 5 52 karyotype (biarmed chromosomes 1, 22, and 23) typical of *P. boylii*, *P. banderanus*, and *P. crinitus*, the karyotype of *P. slevini* differs by having biarmed chromosomes 3 and 9. Compared to the FN 5 58 karyotype (biarmed chromosomes 1, 2, 3, 9, 22, and 23) of *P. mexicanus*-group species, the karyo-

type of *P. slevini* differs by the acrocentric condition of chromosome 2. From the cladistic-based assumption (Rogers et al. 1984; Smith 1990; Stangl and Baker 1984) that the acrocentric condition of chromosome 2 is plesiomorphic for *Peromyscus* and predates the inversions which result in the biarmed conditions of chromosomes 3 and 9, an equal number of inversion events would be needed to explain the differences between the karyotype of *P. slevini* and those of the *P. boylii* and *P. mexicanus* groups, respectively. Cranial similarities of the supraorbital shelf (Carleton 1989), however, support the phylogenetic association of *P. slevini* and the *P. mexicanus*-group as the more likely hypothesis.

The karyotypes of *P. sejugis* from both islands exhibit FN 5 76 but are distinguished by the presence of distal heterochromatin on the short arm of chromo-




Figure 3. Ideogram of various chromosome 13 conditions within the genus *Peromyscus*. The primitive chromosome 13 is the acrocentric condition. The derived biarmed chromosome 13 in *P. maniculatus* (13B) comes from a pericentric inversion between bands A4 and B1. A pericentric inversion between bands A3 and A4 results in the alternative biarmed condition (13B9) found in karyotypes of *P. sejugis* from Isla San Diego (SD). The presence of heterochromatin on the euchromatic short arm of the latter chromosome forms the 13B91 condition indicative of karyotypes of *P. sejugis* from Isla Santa Cruz (not shown).

Isla San Diego provide a character state which would establish *P. sejugis* as a phylogenetic species (see Nixon and Wheeler 1990 and references therein) and support the morphologically based specific distinction of this taxon relative to *P. maniculatus* (Burt 1932). Based on the apparent alternate fixation for 13B9 and 13B91 conditions, a similar argument could be made for a phylogenetic species-based distinction of the two island populations of *P. sejugis*. However, considering the lack of morphologic (Burt 1932), allozymic (Avise et al. 1979) and molecular (Hogan et al. 1997) divergence between these populations we see little value in recommending revision of their current taxonomy.

From the Department of Biology, Texas A&M University, College Station, TX, 77843-3258 (Smith and Greenbaum), and Department of Biology, HQ USAFA/DFB, U.S. Air Force Academy, Colorado Springs, Colorado (Hale). This research was supported by National Institutes of Health, National Institute of General Medical Sciences grant GM 27014 (to I.F.G.) and National Science Foundation grant DEB 9201509 (to D.W.H.). For assistance in the laboratory and/or with collection of the specimens we thank K. M. Hogan, R. R. Hollander, M. Bartlett, S. A. Berend, and S. M. Meyers Unice. S. E. Chirhart, D. M. Deshpande, and J. Weerasinghe provided valuable comments on the manuscript. The animal use in this research was conducted in accordance with the Guide for Care and Use of Laboratory Animals (U.S. Department of Health and Human Services) and approved by the Texas A&M University Animal Care Committee (AUP no. RF91-0250). Address correspondence to I. F. Greenbaum at the address above or e-mail: ira@mail.bio.tamu.edu.

© 2000 The American Genetic Association

References

Allard MW and Greenbaum IF, 1988. Morphological variation and taxonomy of chromosomally differentiated *Peromyscus* from the Pacific Northwest. *Can J Zool* 66:2734-2739.

Allard MW, Gunn SJ, and Greenbaum IF, 1987. Mensural discrimination of chromosomally characterized *Peromyscus oreas* and *Peromyscus maniculatus*. *J Mammal* 68:402-406.

Avise JC, Smith MH, Selander MH, Lawlor TE, and Ramsey PR, 1974. Biochemical polymorphism and systematics in the genus *Peromyscus* V. Insular and mainland species of the subgenus *Haplomylomys*. *Syst Zool* 23: 226-238.

Avise JC, Smith MH, and Selander RK, 1979. Biochemical polymorphism and systematics in the genus *Peromyscus* VII. Geographic differentiation in members of the *truei* and *maniculatus* species groups. *J Mammal* 60: 177-192.

Baker RJ, Haiduk MW, Robbins LW, Cadena A, and Koop BF, 1982. Chromosomal studies of South American bats and their systematic implications. In: *Mammalian biology in South America* (Mares MA and Genoways HH, eds). Special Publication Series, Pymatuning Laboratory of Ecology. Pittsburgh, PA: University of Pittsburgh Press; 303-327.

Burt WH, 1932. Description of heretofore unknown mammals from islands in the Gulf of California, Mexico. *Trans San Diego Soc Nat Hist* 7:161-182.

Burt WH, 1934. Subgeneric allocation of the white-footed mouse, *P. slevini*, from the Gulf of California, Mexico. *J Mammal* 15:159-160.

Calhoun SW and Greenbaum IF, 1991. Evolutionary im-

lications of genic variation among insular populations of *Peromyscus maniculatus* and *Peromyscus oreas*. *J Mammal* 72:248-262.

Carleton MD, 1989. Systematics and evolution. In: *Advances in the study of Peromyscus (Rodentia)* (Kirkland GL Jr and Layne LN, eds). Lubbock, TX: Texas Tech University Press; 7-141.

Greenbaum IF, Gunn SJ, Smith SA, McAllister BF, Hale DW, Baker RJ, Engstrom MD, Hamilton MJ, Modi WS, Robbins LW, Rogers DS, Ward OG, Dawson WD, Elder FFB, Lee MR, Pathak S, Stangl FB Jr, 1994. Cytogenetic nomenclature of deer mice, *Peromyscus* (Rodentia): revision and review of the standardized karyotype. *Cytogenet Cell Genet* 66:181-195.

Gunn SJ, 1988. Chromosomal variation and differentiation among insular populations of *Peromyscus* from the Pacific Northwest. *Can J Zool* 66:2726-2733.

Gunn SJ and Greenbaum IF, 1986. Systematic implications of karyotypic and morphologic variation in mainland *Peromyscus* from the Pacific Northwest. *J Mammal* 67:294-304.

Hogan KM, Davis SK, and Greenbaum IF, 1997. Mitochondrial-DNA analysis of the systematic relationships within the *Peromyscus maniculatus* species group. *J Mammal* 78:733-743.

Hogan KM, Hedin MC, Koh HS, Davis SK, and Greenbaum IF, 1993. Systematic and taxonomic implications of karyotypic, electrophoretic, and mitochondrial-DNA variation in *Peromyscus* from the Pacific Northwest. *J Mammal* 74:819-831.

Hooper ET, 1968. Classification. In: *Biology of Peromyscus (Rodentia)* (King JA, ed). Special publication no. 2. American Society of Mammalogy; 27-74.

Hooper ET and Musser GG, 1964. Notes on the classification of the rodent genus *Peromyscus*. *Occasional Papers of the Museum of Zoology* no. 635. Ann Arbor: University of Michigan Press; 1-13.

Maillaird J, 1924. A new mouse (*Peromyscus slevini*) from the Gulf of California, Mexico. *Proc Calif Acad Sci* 12:1219-1222.

Myers Unice SM, Hale DW, and Greenbaum IF, 1998. Karyotypic variation in populations of deer mice (*Peromyscus maniculatus*) from eastern Canada and the northeastern United States. *Can J Zool* 76:584-588.

Nixon KC and Wheeler QD, 1990. An amplification of the phylogenetic species concept. *Cladistics* 6:211-223.

Osgood WH, 1909. Revision of the mice of the American genus *Peromyscus*. *N Am Fauna* 28:1-285.

Rogers DS, Greenbaum IF, Gunn SJ, and Engstrom MD, 1984. Cytosystematic value of chromosomal inversion data in the genus *Peromyscus* (Rodentia: Cricetidae). *J Mammal* 65:457-465.

Smith SA, 1990. Cytosystematic evidence against monophyly of the *Peromyscus boylii* species group (Rodentia: Cricetidae). *J Mammal* 71:654-667.

Stangl FB Jr, 1986. Aspects of a contact zone between two chromosomal races of *Peromyscus leucopus*. *J Mammal* 67:465-473.

Stangl FB Jr and Baker RJ, 1984. Evolutionary relationships in *Peromyscus*: congruence in chromosomal, genetic, and classical data sets. *J Mammal* 65:643-654.

Sullivan RM, Calhoun SW, and Greenbaum IF, 1990. Geographic variation in genital morphology among insular populations of *Peromyscus oreas*. *J Mammal* 71:48-58.

Sumner AT, 1972. A simple technique for demonstrating centromeric heterochromatin. *Exp Cell Res* 75:304-306.

Verma RS and Babu A, 1995. Banding techniques. In: *Human chromosomes: manual of basic techniques*. Elmsford, NY: Pergamon Press; 74-75.

Received May 26, 1999
Accepted October 24, 1999

Corresponding Editor: Oliver A. Ryder

The Rift Valley Complex as a Barrier to Gene Flow for *Anopheles gambiae* in Kenya: The mtDNA Perspective

T. Lehmann, C. R. Blackston, N. J. Besansky, A. A. Escalante, F. H. Collins, and W. A. Hawley

Descriptions of *A. gambiae* population structure based on microsatellite loci and

Table 1. Polymorphism at the 599 bp segment of the mitochondrial *ND5* gene of *A. gambiae* populations

Collection year	Asembo		Jego	
	1987	1994	1987	1996
<i>N</i> (haplotypes)				
Haplotype diNof				

Table 2. Differentiation between samples from different localities and time points

	Unweighted F_{ST}	Weighted F_{ST}
Asembo 87 vs. Jego 87	20.041 ^{NS}	20.001 ^{NS}
Asembo 94 vs. Jego 96	0.123***	0.176***
Asembo vs. Jego (pooled)	0.089***	0.142***
Asembo 87 vs. Asembo 94	0.026 ^{NS}	0.018 ^{NS}
Jego 87 vs. Jego 96	20.016 ^{NS}	0.005 ^{NS}

*** P . .001.

^{NS} P . .05.

tween localities in the pooled (over time) data. This level of differentiation and derived estimates of gene flow measured by mtDNA, which were adjusted to the difference in N_e between markers, closely agreed with those measured by microsatellites (Lehmann et al. 1998, 1999). The temporal variation between samples taken 7 (Jego) and 9 (Asembo) years apart from the same localities was minimal (Table 2), suggesting that the difference between the original (Besansky et al. 1997) and the present studies were not due to temporal changes in allele frequencies, nor that the 2 years separating the samples from Asembo (1994) and Jego (1996) contributed much to the differentiation between localities. It can be concluded therefore that the discrepancy between the results based on microsatellites and the original mtDNA study was merely a small sample size effect.

The lack of unique alleles in eastern populations and higher F_{ST} than R_{ST} values measured at nine microsatellite loci suggested that pure drift was the main process generating differentiation between these populations (Lehmann et al. 1999). To distinguish between pure drift and mutation-drift using mtDNA data, a test was developed based on the fact that pure drift affects haplotype frequencies but does not systematically affect the number of pairwise substitutions between haplotypes. Accordingly, if two populations became isolated from each other a few generations ago, and one population has experienced a bottleneck and lost several alleles as part of the rapid change in allele frequencies, then allele frequencies will differ markedly between these populations, but the average mutational distance between two different alleles is expected to be the same, regardless of whether they were taken both from a single population or each from a different population. Independent mutations, in addition to drift, must occur in each population to increase the expected mutational distance between two different alleles, each sampled from

Table 3. "Lack-of-fit test" with the mutation-drift model using comparison of haplotype F_{ST} and individual F_{ST} (calculated with weights according to sample sizes)

Populations (sample sizes)	Haplotype F_{ST} (P estimated by permutation test)	Bootstrapped ^b individual F_{ST} (95% CI)
Asembo 94 vs. Jego 96 (25/8)	0.061 (P . .13)	0.171*** (0.072–0.285)
Asembo vs. Jego pooled (32/12)	0.058 (P . .10)	0.138*** (0.070–0.209)
Senegal vs. western Kenya ^a (17/16)	0.046 (P . .029)	0.073*** (0.031–0.134)

*** P . .001.

^a Data from Besansky et al. (1997).

^b Sample sizes were the same as used for the haplotype F_{ST} calculation (shown in first column).

one population. Therefore we calculated F_{ST} on haplotypes instead of individuals (haplotype F_{ST}), which estimates the between-population variation in the number of substitutions (i.e., mutations) per haplotype disregarding the haplotype frequency. If differentiation was generated by pure drift, then the haplotype F_{ST} is expected to be zero. Permutation and bootstrapping tests were used to determine the significance of the results and to evaluate whether an insignificant haplotype F_{ST} reflects low statistical power due to smaller sample size.

The haplotype F_{ST} was calculated between Asembo 1994 and Jego 1996 samples, and in the pooled (over time) samples (Table 3). Haplotype F_{ST} values were approximately one-third of the corresponding individual F_{ST} values and were not significant (P . .09, permutation test), suggesting a lack of fit with the mutation-drift model. To verify that the lack of significance was not a result of weak statistical power due to smaller sample size, we calculated the 95% confidence interval (CI) of individual F_{ST} by bootstrapping over individuals from each population while using the same sample sizes as used for the haplotype F_{ST} calculation. The bootstrapped F_{ST} values were nearly identical to the original individual F_{ST} values (Table 2) and they were significantly higher than zero. Moreover, their lower 95% confidence limits were higher than the corresponding haplotype F_{ST} (Table 3), indicating that the lack of significance of the haplotype F_{ST} values was not due to reduced sample sizes.

In contrast to Kenyan populations across the Rift Valley, higher R_{ST} than F_{ST} values were measured between western Kenya and Senegal (6000 km apart, both west of the Rift Valley barrier) and unique alleles were observed in each population (Lehmann et al. 1996b), suggesting that differentiation between these populations was generated by the mutation-drift model. To test this interpretation, we analyzed the mtDNA data of these populations

(from Besansky et al. 1997). Haplotype F_{ST} for this comparison was significantly larger than zero, and it was not significantly different from the individual F_{ST} based on

From the Division of Parasitic Diseases, Centers for Disease Control and Prevention, Mailstop F22, 4770 Buford Hwy., Chamblee, GA 30341 (Lehmann, Blackstone, Escalante, and Hawley), Department of Biology, Emory University, Atlanta, Georgia (Lehmann), Kenya Medical Research Institute, Clinical Research Centre, Nairobi, Kenya (Hawley), Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana (Besansky and Collins), and Instituto Venezolano de Investigaciones Cientificas, Caracas, Venezuela (Escalante). We are grateful to Mark Danga, Francis Atieli, George Olang, Dr. Charles Mbogo, Mr. Mukoko, and colleagues for assistance in field collections. Dr. Richard R. Hudson reviewed our statistical approach to distinguish pure drift from mutation drift and provided us with valuable comments. Monica Licht, Chuck Porter, Fred Simard, Martin Donnelly, and Fernando Monteiro read earlier versions of this manuscript and provided us with helpful comments. This investigation received financial support from the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (TDR). Address correspondence to T. Lehmann at the address above or e-mail: lbt2@cdc.gov.

References

- Avise JC, 1994. Molecular markers, natural history and evolution. New York: Chapman & Hall.
- Beard CB, Hamm DM, and Collins FH, 1993. The mitochondrial genome of the mosquito *Anopheles gambiae*: DNA sequence, genome organization, and comparisons with mitochondrial sequences of other insects. *Insect Mol Biol* 2:103-124.
- Besansky NJ, Lehmann T, Fahey TG, et al., 1997. Patterns of mitochondrial variation within and between African malaria vectors, *Anopheles gambiae* and *An. arabiensis*, suggests extensive gene flow. *Genetics* 147: 1817-1828.
- Cianchi R, Urbanelli S, Villani F, Sabatini A, and Bullini L, 1985. Electrophoretic studies in mosquitoes: recent



Ovis evolved from a common evolutionary pathway. All arose from ancestral stock

Table 4. Segregation for seed coat texture in different populations of the cross involving IT93K-693-2 and IAR-1696

Population	No. of plants with		χ^2	Probability
	Smooth seeds	Rough seeds		
IT93K-693-2	—	24	—	—
IAR 1696	—	17	—	—
F ₁	26	—	—	—
F ₁ \times IT93K-693-2	9	7	0.25 (1:1)	.5-.7
F ₁ \times IAR 1696	35	32	0.13 (1:1)	.7-.8
F ₂	46	37	0.02 (9:7)	.8-.9

the plants of both parents had rough seed coats, but the F₁ plants had smooth seed coats and brown color, indicating independent gene action for seed coat texture and complete dominance for the brown color (Figure 1). The backcross F₁ population involving IT87D-941-1 segregated into 26 smooth-seeded and 23 rough-seeded plants, and the backcross F₁ population involving Kanannado segregated into 21 smooth-seeded and 24 rough-seeded plants, both fitting closely to a 1:1 ratio. The F₂ population segregated into 138 smooth-seeded and 126 rough-seeded plants, showing close fit to a 9:7 ratio. These data indicate that rough seed coat is controlled by two independent recessive gene pairs, and the recessive gene pair for rough coat in IT89KD-941-1 is different from the gene in Kanannado.

Cross 3: Rough \times Rough

This cross involved a brown-rough-seeded variety, IT93K-693-2, and a white-rough-

←

Figure 2. Seed coat texture and color of (A) IT93K-693-2 (brown-rough), (B) IAR 1696 (white-rough with black hilum), and (C) their F₁ hybrid (black-smooth).



Figure 1. Canine A1AT phenotypes demonstrated by isoelectric focusing and immunoblotting. The bands are the stained immunoprecipitates of alpha 1 antitrypsin. Anode (1) is at the top. The types are 1:M, 2:MS, 3:M, 4:MS, 5:M, 6:S, 7:M, 8:M, 9:S, 10:S. Samples in the following lanes are from the various canides: 1-3: gray wolf, 4-6: domestic dog; 7 and 8: Mexican wolf; 9: coyote; 10: red wolf.

and Pi^S for the common and the slower electrophoretic type, respectively. The phenotypes (band patterns) are called Pi^M , Pi^S , and Pi^{MS} .

Canine A1AT was isolated as described

Table 1. A1AT (Pi) phenotypes

Group	<i>n</i>	M	MS	S
Domestic dog	71	40 (56%)	22 (31%)	9 (13%)
Gray wolf	29	11 (38%)	17 (59%)	1 (3%)
Mexican wolf	20	16 (80%)	4 (20%)	
Red wolf	27			27 (100%)
Coyote	24			24 (100%)
Wolf-dog cross	9	4 (44%)	5 (56%)	

Table 2. Concentrations of A1AT (in mg/ml) \pm standard deviation

Group	Male	Female	Combined
Domestic dog	2.19 \pm 0.38	2.65 \pm 0.42	2.42 \pm 0.41
Gray wolf	2.14 \pm 0.18	2.34 \pm 0.35	2.26 \pm 0.30
Wolf-dog cross	2.28 \pm 0.21	2.45 \pm 0.34	2.37 \pm 0.32
Mexican wolf	2.70 \pm 0.15	3.06 \pm 0.11	2.80 \pm 0.22
Red wolf	2.70 \pm 0.40	3.04 \pm 0.32	2.88 \pm 0.42
Coyote	2.66 \pm 0.40	2.88 \pm 0.38	2.72 \pm 0.43

strated in domestic dogs is also present in gray wolves and that allele frequencies are statistically similar. Mexican wolves are also polymorphic, although the significantly lower Pi^S frequency suggests that they represent a separate population. Red wolves and coyotes are monomorphic for Pi^S . It is as yet undecided if this is due to extensive hybridization or results from common ancestry. Comparison of A1AT concentrations demonstrates the similarity of all animals tested. In addition, the quantitative dimorphism of males and females, known to be present in domestic dogs, was also found in all canids presently tested.

From the Patuxent Wildlife Research Center, Laurel, Maryland (Federoff) and Temple University School of Medicine, Pulmonary Disease Section, 3401 N. Broad St., Philadelphia, PA 19140 (Kueppers). The authors would like to thank the following individuals and institutions for providing blood samples: L. D. Mech (free-ranging Minnesota wolves); F. Knowlton and M. Roetto (Utah coyotes); M. Bush (National Zoo, Washington, DC); N. Reindl (Minnesota Zoo); M. Phillips, V. G. Henry, and J. Gilbreath (U.S. Fish and Wildlife Service red wolf recovery project); W. Waddell and S. Behrns (Point Defiance Zoo and Aquarium, Tacoma, WA); J. Davis (U.S. AWA); S. Johnston and D. Johnston; R. Stubbe; S. Williams and P. Ferrari; S. Lindsey (WCSRC, Eureka, MO); P. Siminski (AZ-Sonora Desert Museum); B. Snyder (Rio Grande Zoo, NM); R. Wack (Columbus Zoo, OH); T. Becker (Zoo America, PA); and S. Fain (U.S.

Fish and Wildlife Service National Forensics Laboratory). N. E. Federoff is currently at the U.S. EPA, 401 M St. SW, Washington, DC.

© 2000 The American Genetic Association

References

- Abrams WR, Kimbal P, and Weinbaum G, 1978. Purification and characterization of canine alpha-1-antitrypsin. *Biochemistry* 17:3556-3561.
- Fagerhol MK and Gedde-Dahl T Jr, 1969. Genetics of the Pi serum types: family studies of the inherited variants of alpha 1 antitrypsin. *Hum Hered* 19:354-359.
- Haldane JBS, 1954. An exact test for the randomness of mating. *J Genet* 52:631-635.
- Hill EP, Sumner PW, and Wooding JB, 1987. Human influences on range expansion of coyotes in the southeast. *Wildl Soc Bull* 15:521-524.
- Hughes D, Elliott DA, Washabau RJ, and Kueppers F, 1995. Effects of age, sex, reproductive status and hospitalization on serum A1AT concentration in dogs. *Am J Vet Res* 56:568-572.
- Koj A, Regoeczi E, Toews CJ, et al., 1978. Synthesis of anti-thrombin III and alpha-1-antitrypsin by the perfused rat liver. *Biochim Biophys Acta* 539:496-504.
- Kueppers F, McConnell IW, and Kramek BA, 1993. Polymorphism of alpha 1 antitrypsin in dogs. *Comp Biochem Physiol* 106B:531-533.
- Li CC, 1955. *Population genetics*. Chicago: University of Chicago Press; pp. 314-327.
- Mancini M, Carbonara AO, and Heremans JF, 1965. Immunochemical quantitation of antigens by single radial immunodiffusion. *Immunochemistry* 2:235-254.
- Muller-Eberhard HJ, 1960. A new supporting medium for preparative electrophoresis. *Scand J Clin Lab Invest* 12:33-37.
- Nowak RM, 1979. *North American quaternary canis*. Monograph no. 6. Lawrence, KS: University of Kansas Museum of Natural History.
- Olsen SJ, 1985. *Origins of the domestic dog: the fossil record*. Tucson, AZ: University of Arizona Press.
- Patterson SD, 1991. Mammalian alpha 1 antitrypsins: comparative biochemistry and genetics of the major plasma serpin. *Comp Biochem Physiol* 100B:439-454.
- Roy MS, Geffen E, Smith D, and Wayne RK, 1996. Molecular genetics of pre-1940 red wolves. *Conserv Biol* 10:1413-1424.
- Travis J and Salvesen GS, 1983. Human plasma proteinase inhibitors. *Annu Rev Biochem* 52:663-674.
- Vila C, Savolainen P, Maldonado JE, Amorim IR, Rice JE, Honeycutt RL, Crandall KA, Lundeberg, and Wayne RK, 1997. Multiple and ancient origins of the domestic dog. *Science* 276:1687-1689.
- Waddell WT and Behrns SK, 1996. *Red wolf 1995 international studbook*. Tacoma, WA: Point Defiance Zoo and Aquarium.
- Wayne RK, 1993. Molecular evolution of the dog family. *Trends Genet* 9:218-224.
- Wayne RK and Jenks SM, 1991. Mitochondrial DNA analysis implying extensive hybridization of the endangered red wolf. *Nature* 351:565-568.
- Wayne RK and Ostrander EA, 1999. Origin, genetic diversity and genome structure of the domestic dog. *BioEssays* 21:247-257.

Received September 1, 1998

Accepted October 24, 1999

Corresponding Editor: Robert Wayne