

Phylogeographical pattern correlates with Pliocene mountain building in the alpine scree weta (Orthoptera, Anostostomatidae)

S. A. TREWICK, G. P. WALLIS and M. MORGAN-RICHARDS

Department of Zoology, University of Otago, PO Box 56, Dunedin, New Zealand

Abstract

Most research on the biological effects of Pleistocene glaciation and refugia has been undertaken in the northern hemisphere and focuses on lowland taxa. Using single-strand conformation polymorphism (SSCP) analysis and sequencing of mitochondrial cytochrome oxidase I, we explored the intraspecific phylogeography of a flightless orthopteran (the alpine scree weta, *Deinacrida connectens*) that is adapted to the alpine zone of South Island, New Zealand. We found that several mountain ranges and regions had their own reciprocally monophyletic, deeply differentiated lineages. Corrected genetic distance among lineages was 8.4% (Kimura 2-parameter [K2P]) / 13% (GTR + I + Γ), whereas within-lineage distances were only 2.8% (K2P) / 3.2% (GTR + I + Γ). We propose a model to explain this phylogeographical structure, which links the radiation of *D. connectens* to Pliocene mountain building, and maintenance of this structure through the combined effects of mountain-top isolation during Pleistocene interglacials and ice barriers to dispersal during glacials.

Keywords: alpine, COI, mitochondrial DNA, New Zealand, phylogeography, Pleistocene glaciation

Received 23 September 1999; revision received 12 November 1999; accepted 12 November 1999

Introduction

Biologists live and work in what is in effect an interglacial. The 10 000 years or so since the end of the last glacial is a period similar in duration to each of the 20 or so interglacials of the Pleistocene (Stevens 1981; Cooper & Millener 1993). This brings an immediacy to the study of the evolutionary effects of the Pleistocene glacial epoch, which has stimulated considerable research interest in the field. Studies of the impact of such widespread and recent events reveal not only the nature of the world today but provide an insight into more ancient evolutionary processes.

Although the physical effects of the Pleistocene glacials were global, they were expressed in different ways regionally (Webb & Bartlein 1992; Cox & Moore 1993). Biological response to glacial–interglacial cycling appears to vary with locality and taxa (Jaarola & Tegelström 1995;

Hewitt 1996; Zink 1996; Roy 1997; Taberlet *et al.* 1998; and references therein). Most studies of the biological effects of glaciation have been concerned with northern hemisphere regions where the last glacial epoch involved southward extension of continuous ice sheets. This ice erased biota from many regions of northern Europe and North America and forced southward range movement as climate patterns shifted (Comes & Kadereit 1998). In western Europe, for instance, several studies have indicated the existence of refugia where the biota became isolated between the advancing ice front and the Mediterranean Sea (see for examples: Comes & Kadereit 1998; Lunt *et al.* 1998; Taberlet *et al.* 1998). The final recession of ice sheets (\approx 14 000 years ago) opened up extensive areas that were recolonized from the refugia. The phylogeographical effects of Pleistocene glaciation were, however, more complex owing to the cycling of multiple glacial and interglacial episodes.

In the southern hemisphere, the distance of the southern continents from the south pole and the circumpolar current prevented extension of ice sheets from Antarctica to other

Correspondence: Steven A. Trewick. Fax: +64 3 479 7584; E-mail: steve.trewick@stonebow.otago.ac.nz

subantarctic land-bridging (Fleming 1963, 1979; Wardle 1963). The alpine invertebrate fauna is diverse, highly endemic and specialized (Mark & Dickinson 1997), and several groups show well-developed freeze tolerance (Block *et al.* 1998; Sinclair *et al.* 1999). The evidence of the impact of the Pleistocene glacial epoch, both physical (e.g. glacier erosion surfaces, moraines, valleys and fjords) (Stevens 1981; Pillans *et al.* 1992) and biological (extinction, zonation, alpine specialization) is abundant (Willet 1950; Fleming 1963; Burrows 1965; Dumbleton 1970; McGlone 1985, 1988). Fleming (1979) proposed a Pleistocene glaciation model of forced adaptation for speciation in New Zealand parrots (*Nestor*) and suggested that this would be a 'plausible mechanism for the occupation of the alpine zone by many plants and insects (e.g. *Hemideina* and *Deinacrida*)'. Yet to date there have been few phylogeographical studies exploring the molecular evidence for the biological effects of these events (but see Emerson & Wallis 1995; King *et al.* 1996; Buckley *et al.* 1998).

We have used a molecular approach to examine the intraspecific phylogeography of one of New Zealand's largest and most notable alpine invertebrates, the scree weta *Deinacrida connectens* (Orthoptera: Anostostomatidae). Scree weta are widespread in South Island but absent from North Island. The species inhabits alpine scree slopes 1200–3600 m above sea level and is limited to this zone by as yet undetermined factors. The range cut-off at the lower altitudinal limit on each mountain is not consistent with any outwardly apparent features and exists despite the extension of apparently suitable scree habitat below this altitude. The scree weta appears to survive freezing without ill effects (B. Sinclair, personal communication). The physiological restriction to high altitude and flightlessness of this insect mean that in current climatic conditions, populations can be isolated from one another, even on mountain tops within the same ranges.

Studies of the processes governing the geographical distributions of genealogical lineages (Riddle 1996) are particularly well served in the intraspecific context by the use of mitochondrial DNA (mtDNA) analysis (Avise *et al.* 1987; Avise 1994). The lack of recombination and maternal inheritance allow historical information to be retained intact even where a complex glacial history may have fostered repeated dispersal events that lead to blending of the nuclear genome through recombination.

We have two alternative broad hypotheses relating to various possible patterns that we might find. We refer to these as: (i) alpine radiation and (ii) recent isolation models. The alpine radiation model proposes that the origin of the species and its special adaptations coincided with major alpine uplift and onset of the Pleistocene glacial epoch. If this model is correct, then coalescence should fall within the time of initial alpine uplift (2–5 Ma), with marked phylogeographical structuring. Adaptation

to the alpine environment could have occurred either in parallel on the different ranges, or at an early stage in one location with subsequent spread to multiple emerging ranges. Alternatively, our recent isolation model proposes that all populations were effectively in contact until either: (i) modern populations became isolated in alpine regions as the climate warmed at the end of the Pleistocene (< 14 000 years ago); or (ii) habitat modification and introduction of mammalian predators (e.g. rats, Holdaway 1996) within the last 2000 years led to a more fragmented distribution. Under these circumstances, coalescence time would depend on effective population size, but would probably be quite recent, and there should be little, if any, phylogeographical structure. Population isolation as a result of predation seems less probable because the species appears to be physiologically restricted to altitudes well above the point at which predators are thought to impact (Gibbs & Richards 1990).

Thus, the timings of our two alternative hypotheses differ by more than two orders of magnitude, and should be distinguishable using the methods we propose. In fact, genetic differentiation will be minimal on a timescale of at least an order of magnitude greater than that predicted by our recent isolation models, so cessation of pre-existing gene flow at any time during the Holocene would probably be undetected.

Materials and methods

Sampling

Specimens of *Deinacrida connectens* were collected from alpine scree slopes throughout South Island, New Zealand, in 1991 (Morgan-Richards & Gibbs 1996) and 1998 (Fig. 1). Populations at the following locations were sampled (name with abbreviation underlined, longitude/latitude, sample size): Mt Peel, 41°08'S 172°35'E, $n = 7$; Mt Arthur, 41°13'S 172°42'E, $n = 8$; Spence Peak, 44°52'S 167°51'E, n

5 μ L of 10 mg/mL proteinase K in 600 μ L of TNES buffer (20 mM EDTA, 50 mM Tris, 400 mM NaCl, 0.5% sodium dodecyl sulphate [SDS]) at 50 °C. Ten per cent 5 M NaCl was added and the extractions were shaken vigorously for 20 s followed by centrifugation at 16 060 g for 5 min. The supernatant was removed and precipitated with an equal volume of cold 100% ethanol. DNA was collected by centrifugation and washed with 70% ethanol before being dried and resuspended in water.

Our molecular analysis used primers that target regions of the mtDNA cytochrome oxidase I gene (COI). These primers are known to be highly conserved and applicable to a wide range of invertebrate taxa (Lunt *et al.* 1996), and the COI gene has been successfully utilized in intra- and interspecific studies (Zhang & Hewitt 1996; Funk 1999).

Single-stranded conformation polymorphism analysis

Mitochondrial primers C1-J-1718 and C1-N-2191 (Simon *et al.* 1994) were used to amplify a short (\approx 400 bp) fragment of the 5' end of the COI. Single-stranded conformation polymorphism (SSCP) analysis was used to screen for variant haplotypes. Polymerase chain reaction (PCR) products were isotopically labelled by incorporation of [α^{33} P]-dATP. Ten-microlitre reaction mixtures (200 μ M dNTPs, 2.5 mM MgCl₂, 0.25 U of Qiagen *Taq*) were amplified as follows: an initial denaturation at 94 °C for 60 s followed by 40 cycles of 94 °

and predator introduction). The estimated mean coalescence time of ~ 4 Myr falls within the time range estimated for the emergence of the main axial range of South Island mountains, and the polytomous str

would presumably have forced the geographical range of alpine organisms to fluctuate accordingly. Two features of glaciation will, however, have had subtly different effects. A colder climate will have lowered the alpine zone and allowed *D. connectens* to colonize lower altitudes with potentially greater habitat area. At the same time, however, glacier extension would have prevented colonization of many valley systems and maintained lowland barriers between scree populations (Fig. 3). Populations in the vicinity of Eyre Peak and Rees Saddle, for instance, would have remained separated by ice (see Fig. 1 inset). In areas where ice was sparse, such as Marlborough, exchange between populations by dispersal across valleys would have been feasible. The presence of glaciers at the northern end of the Southern Alps may have prevented gene flow between weta populations in Marlborough and Nelson, as has been proposed for alpine grasshoppers in the region (Peterson 1968). It would certainly have limited exchange between Nelson and the other populations to the south. Warm interglacials will have restricted the range of *D. connectens* to higher altitudes and may even have led to local extinction.

Although the phylogeographical structure of mtDNA haplotypes is very clear, it is only vaguely paralleled by morphological and chromosome variation among populations. Colour variation shows some north-to-south clustering (Field 1980), but colour variation within regions such as Marlborough (lineage A) is high (Morgan-Richards & Gibbs 1996). Southern populations had karyotype numbers (XO males) of 17 or 19, whereas those from Fox Peak northwards had 21 (Morgan-Richards & Gibbs 1996). There is good bootstrap support (85%) for a split between the mtDNA lineages (D, F, G vs. A, B, C, E) that coincides with this karyotypic differentiation (Fig. 2). Within these two (north and south) groups, the distribution of other, more subtle, karyotype variation appears to overlay mtDNA structure. Notably, weta on Mt Arthur (lineage C) have a similar karyotype to populations in Marlborough (lineage A), and at St Mary's Range haplotype F2 was found in an individual with 17 chromosomes whilst the same haplotype was present among individuals with 19 chromosomes at Rees Saddle. This level of intra-specific karyotype variation is a feature of Orthoptera (White 1978) and typical of the family (Morgan-Richards 1997).

Quaternary biogeography is dominated by postglacial range expansion from glacial age refugia that were positioned away from cold and arid areas (e.g. Cwynar & MacDonald 1987; Bennett *et al.* 1991; Hewitt 1993; Cooper *et al.* 1995; Joseph *et al.*

- within New Zealand and their ecological significance. Part II. *Tuatara*, **13**, 9–29.
- Comes HP, Kadereit JW (1998) The effect of Quaternary climatic changes on plant distribution and evolution. *Trends in Plant Science*, **3**, 432–438.
- Cooper RA, Millener PR (1993) The New Zealand biota: historical background and new research. *Trends in Ecology and Evolution*, **8**, 429–433.
- Cooper SJB, Ibrahim KM, Hewitt GM (1995) Postglacial expansion and subdivision of the grasshopper *Chorthippus parallelus* in Europe. *Molecular Ecology*, **4**, 49–60.
- Cox CB, Moore PD (1993) *Biogeography: an ecological and evolutionary approach*, 5th edn. Blackwell Scientific Publications, Oxford.
- Cwynar LC, MacDonald GM (1987) Geographical variation of lodgepole pine in relation to population history. *American Naturalist*, **129**, 463–469.
- Denton GH, Hughes TJ (1981) *The Last Great Ice Sheets*. Wiley, New York.
- Dumbleton LJ (1970) Pleistocene climates and insect distributions. *New Zealand Entomologist*, **4**, 3–23.
- Emerson BC, Wallis GP (1995) Phylogenetic relationships of the *Prodontria* (Coleoptera: Scarabaeidae; Subfamily Melolonthinae), derived from sequence variation in the mitochondrial cytochrome oxidase II gene. *Molecular Phylogenetics and Evolution*, **4**, 433–447.
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distance among DNA haplotypes: applications to human mitochondrial DNA restriction data. *Genetics*, **131**, 479–491.
- Field LH (1980) Observations on the biology of *Deinacrida connectens* (Orthoptera: Stenopematidae), an alpine weta. *New Zealand Journal of Zoology*, **7**, 211–220.
- Fleming CA (1963) The age of the alpine biota. *Proceedings of the New Zealand Ecological Society*, **10**, 15–18.
- Fleming CA (1979) *The Geological History of New Zealand and its Life*. University of Auckland and Oxford University Press, Auckland.
- Flint RF (1957) *Glacial and Pleistocene Geology*. John Wiley & Sons, New York.
- Funk DJ (1999) Molecular systematics of cytochrome oxidase I and 16S from *Neochlamisus* leaf beetles and the importance of sampling.

- model. *Proceedings of the Royal Society London B*, **264**, 1337–1344.
- Schneider S, Kueffer J-M, Roessli D, Excoffier L (1997) *ARLEQUIN*, Version 1.1: software for populations genetics data analysis. Genetics and Biometry Laboratory, University of Geneva, Switzerland.
- Simon C, Frati F, Beckenbach A *et al.* (1994) Evolution, weighting and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the American Entomological Society*, **87**, 651–701.
- Sinclair BJ, Worland MR, Wharton DA (1999) Ice nucleation and freezing tolerance in New Zealand alpine and lowland weta, *Hemideina* spp. (Orthoptera: Stenopelmatidae). *Physiological Entomology*, **24**, 56–63.
- Stevens G (1981) *New Zealand Adrift: the Theory of Continental Drift in a New Zealand Setting*, 2nd edn. Reed, Wellington.
- Sunnucks P, Hales DF (1996) Numerous transposed sequences of mitochondrial cytochrome oxidase I–II in aphids of the genus *Sitobion* (Hemiptera: aphididae).